

Chapter 24 From Petroleum to Pharmaceuticals

24.1 Petroleum Refining and the Hydrocarbons

24.2 Functional Groups and Organic Synthesis

24.3 Pesticides and Pharmaceuticals

Physical Properties of Alkanes and Cycloalkanes

Why do molecules tend to stick together?

Boiling Points


increase with increasing number of carbons

more atoms, more electrons, more opportunities for induced dipole-induced dipole forces

decrease with chain branching

branched molecules are more compact with smaller surface area — fewer points of contact with other molecules

Boiling points and melting points of n-alkanes



Boiling Points

increase with increasing number of carbons

more atoms, more electrons, more
opportunities for induced dipole-induced
dipole forces



Heptane
bp 98°C



Octane
bp 125°C



Nonane
bp 150°C

Boiling Points

decrease with chain branching (WHY?)

branched molecules are more compact with smaller surface area—fewer points of contact with other molecules



n-octane: bp 125°C



2-Methylheptane: bp 118°C



2,2,3,3-Tetramethylbutane: bp 107°C

Boiling Points of Alkanes

governed by strength of intermolecular attractive forces

alkanes are nonpolar, so dipole-dipole and dipole-induced dipole forces are absent

only forces of intermolecular attraction are induced dipole-induced dipole forces

Induced dipole-Induced dipole attractive forces



two nonpolar molecules

center of positive charge and center of negative charge coincide in each

Induced dipole-Induced dipole attractive forces



movement of electrons creates an
instantaneous dipole in one molecule (left)

Induced dipole-Induced dipole attractive forces



temporary dipole in one molecule (left)
induces a complementary dipole in other
molecule (right)

Induced dipole-Induced dipole attractive forces



temporary dipole in one molecule (left)
induces a complementary dipole in other
molecule (right)

Induced dipole-Induced dipole attractive forces



the result is a small attractive force between the two molecules

Induced dipole-Induced dipole attractive forces




the result is a small attractive force between the two molecules

Straight chain hydrocarbon

Branched hydrocarbon

Lots of intermolecular contacts

Fewer intermolecular contacts



Small organic molecules: required to sustain life and health

Life: food (pesticides and herbicides)


Health: disease and pain control (pharmaceuticals)

Large organic molecules: Polymers (many "mers" or repeating units)


Biomolecules: the blue print (DNA) and the machinery (proteins and sugars) of life

Organic short hand for structures

Note hydrocarbon skeleton, recognize functional groups




Isoamyl acetate




Benzyl acetate

Structure of some small molecule analgesics


Molecular shape and functional groups determine the onset of pain and the molecular structure for its relief



[a] Aspirin




[b] Acetaminophen
(4-acetaminophenol)




[c] Morphine

Structure of some small molecule pesticides


Kill insects indiscriminately. Also be toxic to humans.




[a] Nicotine



[b] DDT




[c] Malathion




[d] Methoprene

Structure of some small molecule herbicides


Kill only insects; some attract insects and make them sterile



[a] 2,4-D



[b] TCDD




[c] Atrazine

Structure of some small molecule antibiotics and


Bacteria mistake molecule (a) for a molecule they need to make folic acid. This mistake kills them.



[a] Sulfanilamide



[b] Penicillin G




[c] Tetracycline


Structure of some steroids

Cholesterol is found in all tissues of the body


Human sex hormones are structurally based on cholesterol



[a] Cholesterol



[b] Cortisone



R = —OH (testosterone)

R = —C(=O)CH₃ (progesterone)

[c] Progesterone and testosterone