

18.1

Electron Waves and  
Chemical Bonds

## *Models for Chemical Bonding*


The Lewis model of chemical bonding predates the idea that electrons have wave properties.

There are two other widely used theories of bonding that are based on the wave nature of an electron.

Valence Bond Theory

Molecular Orbital Theory

*Consider formation of  $H_2$  from two hydrogen atoms.*



Examine how the electrostatic forces change as two hydrogen atoms are brought together.

These electrostatic forces are:

attractions between the electrons and the nuclei

repulsions between the two nuclei

repulsions between the two electrons

Figure 1.14




Figure 1.14




Figure 1.14





Figure 1.14




2 H atoms: each electron "feels"  
attractive force of one proton



$H_2$  molecule: each electron "feels"  
attractive force of both protons

Figure 1.14



## *Models for Chemical Bonding*

### Valence Bond Theory

constructive interference between two electron waves is basis of shared-electron bond

### Molecular Orbital Theory

derive wave functions of molecules by combining wave functions of atoms

18.1  
Bonding in H<sub>2</sub>:  
The Valence Bond Model


## *Valence Bond Model*

Electron pair can be shared when half-filled orbital of one atom overlaps in phase with half-filled orbital of another.

## Valence bond model



*in-phase overlap of two half-filled  
hydrogen 1s orbitals*




$\sigma$  bond of  $\text{H}_2$

## *Valence Bond Model*

$\sigma$  Bond: orbitals overlap along internuclear axis

Cross section of orbital perpendicular to internuclear axis is a circle.



## *Valence Bond Model of $H_2$*




Figure 1.17(a) The 1s orbitals of two separated hydrogen atoms are far apart. Essentially no interaction. Each electron is associated with a single proton.

## *Valence Bond Model of $H_2$*




Figure 1.17(b) As the hydrogen atoms approach each other, their 1s orbitals begin to overlap and each electron begins to feel the attractive force of both protons.

## *Valence Bond Model of H<sub>2</sub>*




Figure 1.17(c) The hydrogen atoms are close enough so that appreciable overlap of the the two 1s orbitals occurs. The concentration of electron density in the region between the two protons is more readily apparent.

## *Valence Bond Model of $H_2$*




Figure 1.17(d) A molecule of  $H_2$ . The two hydrogen 1s orbitals have been replaced by a new orbital that encompasses both hydrogens and contains both electrons.

18.1  
Bonding in H<sub>2</sub>:  
The Molecular Orbital Model

## *Main Ideas*

Electrons in a molecule occupy molecular orbitals (MOs) just as electrons in an atom occupy atomic orbitals (AOs).

Two electrons per MO, just as two electrons per AO.

Express MOs as combinations of AOs.

## MO Picture of bonding in $H_2$

Linear combination of atomic orbitals method expresses wave functions of molecular orbitals as sums and differences of wave functions of atomic orbitals.

Two AOs yield two MOs

Bonding combination

Antibonding combination

$$\psi_{MO} = \psi(H)_{1s} + \psi(H')_{1s} \quad \psi'_{MO} = \psi(H)_{1s} - \psi(H')_{1s}$$

*Fig. 1.19 Energy-level diagram for H<sub>2</sub> MOs*

1s —  
AO

— 1s  
AO

*Fig. 1.19 Energy-level diagram for H<sub>2</sub> MOs*

MO

—  $\sigma^*$  antibonding

—  $\sigma$  bonding

MO

*Fig. 1.19 Energy-level diagram for H<sub>2</sub> MOs*




Fig. 1.19 Energy-level diagram for  $H_2$  MOs

