


# Entodermal derivatives: formation of the gut, liver, and pancreas

Mike Gershon



The background of the slide is a dark blue gradient. At the bottom, there are several faint, concentric circular patterns that resemble ripples in water, centered horizontally and slightly offset vertically.

# Folding forms the gut

- Primitive gut extends from buccopharyngeal to cloacal membrane.
  - Move toward each other
- Cardiogenic mesenchyme is originally rostral, but folding brings it caudal to buccal membrane.
- Foregut and hindgut become recognizable
- Portion of yolk sac is incorporated into the embryo as bowel.
- Midgut remains open.





# Cephalocaudal and lateral folding occur simultaneously



- Meeting and fusion of cranial, lateral, and caudal edges of the embryo create the primordial foregut and hindgut
  - Slow fusion of midgut-due to presence of yolk sac. Midgut remains open until week 6-connects to yolk sac via *vitelline duct*.
  - Buccopharyngeal membrane opens at 4 and cloacal membrane at 7 weeks

# Flexion delimits the bowel



- After the gut forms, it is attached to the body wall by dorsal and ventral mesenteries; ventral is lost except in region of liver. Vertebral duct remains in umbilical cord.

IN CROSS SECTION




Fig. 3.

c) The ventral wall begins to form (fig. 3). The gut is delineated, surrounded by intra-embryonic coelom.

d) At the level of the umbilicus, the ventral wall does not close completely (fig. 4).






Fig. 4.

# Anterior-posterior and lateral folding form the primitive gut



- Embryonic disc grows faster in length than the yolk sac causing the embryo to bend.
  - Dorsal surface grows more rapidly than the ventral
- Lateral folding
  - Fusion with apposing side except in the region of the yolk sac, and allantois
- Folding brings the heart and septum transversum caudal to buccopharyngeal membrane.



**Figure 6–3.** Formation of the dorsal mesentery. *A*, The primitive gut tube initially hangs from the posterior body wall by a broad bar of mesenchyme but, *B*, in regions inferior to the septum transversum this connection thins out to form a membranous dorsal mesentery composed of reflected peritoneum.

The dorsal mesentery thins to allow the gut to be flexibly suspended


# The foregut has many derivatives

- Pharynx and its derivatives
- Lower Respiratory tract
- Esophagus
- Stomach
- Duodenum proximal to ampulla of Vater
- Liver
- Biliary Apparatus
- Pancreas


**From stomach to biliary apparatus, all are supplied by the celiac artery, “the artery of the foregut.”**

# Esophagus elongates rapidly

- Appears to grow faster at its cranial than caudal end.
- Stomach does not descend but arises from a region just caudal to septum transversum that has been fated to be stomach.
- Epithelium obliterates lumen of esophagus and is recanalized by apoptosis (week 8).
  - Failure causes polyhydramnios
  - Esophageal atresia or tracheo-esophageal fistula.
- Stomach enlarges and rotates




# Obliteration of the lumen and recanalization occurs



**Figure 9-13.** Formation of the definitive gut lumen. Proliferation of the endodermal lining completely occludes the gut tube during the sixth week. Recanalization is completed by week 9. Incomplete or abnormal recanalization may result in duplication of the lumen or stenosis of the gut tube.


# The stomach rotates 90° in a clockwise direction




- Dorsal surface grows faster than the ventral to create the greater and lesser curvature. Acquires a transverse position

# Rotation of the stomach creates the lesser sac


- Dorsal mesogastrium moves to left.
- Ventral mesogastrium attaches to liver and body wall.
- Inferior recess form the greater omentum
  - Layers fuse to obliterate the lesser sac



# Rotation of the stomach forms the omental bursa




# Movements of the mesentery and stomach are made possible by vacuolization due to selective apoptosis




**Figure 9-4.** The rotation of the stomach around its longitudinal axis commences with vacuolization of the right side of the thick mesenchymal bar that initially suspends the stomach from the posterior body wall.

# Liver, biliary system and pancreas arise from the duodenum




**Figure 9-6.** Development of the liver, gallbladder, pancreas, and their duct systems from endodermal diverticula of the duodenum. The liver bud sprouts during the fourth week and expands in the ventral mesentery. The cystic diverticulum and ventral pancreatic bud also grow into the ventral mesentery, whereas the dorsal pancreatic bud grows into the dorsal mesentery. During the fifth week, the ventral pancreatic bud migrates around the posterior side (former right side) of the duodenum to fuse with the dorsal pancreatic bud. The main duct of the ventral bud ultimately becomes the major pancreatic duct, which drains the entire pancreas.

# Hepatic diverticulum grows from the duodenum into the ventral mesentery




- Begins ~ week 4
- Divides into cranial and caudal buds.
- **Cranial bud grows faster and becomes the hepatic parenchyma;**
  - Hematopoietic colonists arrive ~ week 6
- **Caudal bud gives rise to the biliary system.**

# Ventral mesentery forms falciform ligament, hepatic peritoneum, and lesser omentum




**Figure 9–8.** Formation of the liver and associated membranes. As the liver bud grows into the ventral mesentery, its expanding crown makes direct contact with the developing diaphragm. The ventral mesentery that encloses the growing liver bud differentiates into the visceral peritoneum of the liver, which is reflected onto the diaphragm. This zone of reflection, which encircles the area where the liver directly contacts the diaphragm (the bare area), becomes the coronary ligament. The remnant of ventral mesentery connecting the liver with the anterior body wall becomes the falciform ligament, whereas the ventral mesentery between the liver and lesser curvature of the stomach forms the lesser omentum.

# Ventral mesogastrium supports liver and stomach




# Rotation of the stomach shapes the pancreas




- **Pancreas arises from dorsal and ventral buds.**
- **Rotation brings ventral to dorsal bud.**
- **Buds fuse.**
  - **Ventral duct becomes the main pancreatic duct but the dorsal bud forms most of the pancreas**
  - **Ventral bud forms only the uncinete process and inferior part of the head of the pancreas.**

# Aberrant rotation causes an annular pancreas



**Figure 9-7.** The ventral pancreas may consist of two lobes. If the lobes migrate around the duodenum in opposite directions to fuse with the dorsal pancreatic bud, an annular pancreas is formed.



**Figure 9-2.** Structure of the gut tube. The foregut consists of the pharynx, located superior to the respiratory diverticulum, the thoracic esophagus, and the abdominal foregut. The abdominal foregut forms the abdominal esophagus, stomach, and about half of the duodenum and gives rise to the liver, the gallbladder, the pancreas, and their associated ducts. The midgut forms half of the duodenum, the jejunum and ileum, the ascending colon, and about two thirds of the transverse colon. The hindgut forms one third of the transverse colon, the descending colon and sigmoid colon, and the upper two thirds of the anorectal canal. The abdominal esophagus, stomach, and superior part of the duodenum are suspended by dorsal and ventral mesenteries; the abdominal gut tube excluding the rectum is suspended in the abdominal cavity by a dorsal mesentery only.

# Derivatives of the midgut

- Small intestine (except for the proximal duodenum).
- Cecum
- Appendix
- Ascending colon
- Right 1/2 to 2/3 of the proximal transverse colon
- All are supplied by the superior mesenteric artery (“the artery of the midgut”)

# The midgut grows rapidly and herniates into the umbilical cord






Figure 9-5. Development of the greater omentum and lesser sac. A, B, The rotation of the stomach and growth of the dorsal mesogastrium create a sac (the greater omentum) that dangles from the greater curvature of the stomach. B, C, When the duodenum swings to the right, it becomes secondarily fused to the body wall, enclosing the space posterior to the stomach and within the expanding cavity of the greater omentum. This space is the lesser sac of the peritoneal cavity. The remainder of the peritoneal cavity is now called the greater sac. The principal passageway between the greater and lesser sacs is the epiploic foramen of Winslow.

Week 6





**Figure 9-9.** Hemiation and rotations of the intestine. **A, B,** At the end of the sixth week, the primary intestinal loop herniates into the umbilicus, rotating through 90 degrees counterclockwise (in frontal view). **C,** The small intestine elongates to form jejunal-ileal loops, the cecum and appendix grow, and, at the end of the 10th week, the primary intestinal loop retracts into the abdominal cavity, rotating an additional 180 degrees counterclockwise. **D, E,** During the 11th week, the retracting midgut completes this rotation as the cecum is positioned just inferior to the liver. The cecum is then displaced inferiorly, pulling down the proximal hindgut to form the ascending colon. The descending colon is simultaneously fixed on the left side of the posterior abdominal wall. The jejunum, ileum, and transverse colon and sigmoid colon remain suspended by mesentery.

The midgut rotates around an axis of the superior mesenteric artery:


1. 90°
2. 180°


Midgut hernia reduced at week 10.

# Rotation of the midgut

- 1. Cranial and caudal loop form.
- 2. Cranial growth >>> caudal growth.
- 3. Apex of loop is vitelline duct.
- 4. Cranial loop moves to right and caudal loop to left (90° counterclockwise).
- 4. Reduction of midgut hernia with rotation a further 180°.
  - Brings cecum to right
  - Moves down
  - Becomes secondarily retroperitoneal.


# Loops of bowel fuse with the body wall and become secondarily retroperitoneal





**Figure 9-10.** Intraperitoneal, retroperitoneal, and secondarily retroperitoneal organs of the abdominal gastrointestinal tract.


# Volvulus is a serious complication of excessive flexibility




**Figure 9-20.** Volvulus. Volvulus may occur as suspended regions of the gut twist around themselves, constricting the intestine and/or compromising its blood supply.




**A** Persistent attachment to umbilicus      Meckel's diverticulum



**B** Omphalomesenteric fistula



**C** Omphalomesenteric cyst




**D** Omphalomesenteric ligament (fibrous band)

**Figure 9-21.** Meckel's diverticulum. *A*, A typical Meckel's diverticulum is a finger-like projection of the ileum located about 100 cm proximal to the cecum. A Meckel's diverticulum may form (*B*) a patent fistula connecting the umbilicus with the ileum, (*C*) an isolated cyst suspended by ligaments, or (*D*) a fibrous band connecting the ileum and anterior body wall at the level of the umbilicus. (Drawings courtesy of Children's Hospital Medical Center, Cincinnati, OH.)


# Derivatives of the hindgut

- Left 1/3 to 1/2 of the distal transverse colon
- Descending colon
- Sigmoid colon
- Rectum
- Superior part of anal canal
- Epithelium of urinary bladder and most of the urethra
- **All are supplied by the inferior mesenteric artery, “the artery of the”. hindgut**


# The hindgut is originally a cloaca-partioned to form rectum and urogenital sinus



# Urorectal septum divides the cloaca




**Figure 9–11.** Subdivision of the cloaca into an anterior primitive urogenital sinus and a posterior rectum between 4 and 6 weeks. The urorectal septum that divides the cloaca is composed of three distinct septa. Initially, a superior Tourneux fold grows inferiorly to the level of the future pelvic urethra. Separation is then completed by left and right Rathke folds that grow in a coronal plane.



**Figure 9–12.** The lower third of the anorectal canal is formed by an ectodermal invagination called the anal pit. The border between the superior end of the anal pit and the inferior end of the rectum is demarcated by mucosal folds called the pectinate line in the adult.

**Hindgut  
forms  
superior 2/3  
of rectal  
canal;  
proctodeum  
forms lower  
1/3; divided  
at pectinate  
line**

Never forget  
the pectinate  
line



# If anything can go wrong it will; anorectal malformations




Fig. 2. — Imperforate anus with scrotal fistula.




Fig. 3. — Rectal atresia.




Fig. 4. — Rectal atresia with urinary fistula.






Fig. 5. — Rectal atresia with vaginal fistula.

# The END


Have a nice day!





**Timeline.** Development of the gut tube and its derivatives.





**Figure 9-22.** Fate of the allantois. *A*, Normally, the allantois becomes occluded to form the urachus or median umbilical ligament of the adult. Very rarely, parts of the allantois may remain patent, producing (*B*) a urachal fistula, (*C*) a urachal sinus, (*D*) a urachal diverticulum, or (*E*) a urachal cyst.



Table 9-2

## Derivatives of the Septum Transversum

REGIONS OF SEPTUM TRANSVERSUM	DERIVATIVES
Cranial region	Central tendon of the diaphragm Myocytes of the pleuroperitoneal membranes
Central mesenchyme	Hematopoietic cells of liver
Caudal region (ventral mesentery)	Falciform ligament Visceral peritoneum of the liver, including the coronary ligament Visceral peritoneum of the gallbladder Lesser omentum, including the hepatoduodenal and hepatogastric ligaments

**Table 9-1**  
**The Derivatives of the Primitive Gut Tube**

REGIONS OF THE DIFFERENTIATED GUT TUBE	ACCESSORY ORGANS DERIVED FROM THE GUT TUBE ENDODERM
<i>Foregut</i>	
Pharynx	Pharyngeal pouch derivatives (see Ch. 12)
Thoracic esophagus	Lungs
Abdominal esophagus	
Stomach	
Superior half of duodenum (superior to the ampulla of Vater)	Liver parenchyma and hepatic duct epithelium Gallbladder, cystic duct, and common bile duct Dorsal and ventral pancreatic buds (exocrine cells and pancreatic duct epithelium; probably also pancreatic endocrine cells)
<i>Midgut</i>	
Inferior half of duodenum	
Jejunum	
Ileum	
Cecum	
Appendix	
Ascending colon	
Right two thirds of transverse colon	
<i>Hindgut</i>	
Left one third of transverse colon	
Descending colon	
Sigmoid colon	
Rectum	Urogenital sinus and derivatives (see Ch. 10)



**Figure 9-1.** A, The foregut, midgut, and hindgut of the primitive gut tube are formed by the combined action of differential growth and lateral and cephalocaudal folding. The foregut and hindgut are blind-ending tubes that terminate at the buccopharyngeal and cloacal membranes, respectively. The midgut is at first completely open to the cavity of the yolk sac B, C, As folding proceeds, however, this connection is constricted to form the narrow vitelline duct (D).