



Entodermal derivatives: formation of the gut, liver, and pancreas

Mike Gershon

Folding forms the gut



- Primitive gut extends from buccopharyngeal to cloacal membrane.
 - Move toward each other
- Cardiogenic mesenchyme is originally rostral, but folding brings it caudal to buccal membrane.
- Foregut and hindgut become recognizable
- Portion of yolk sac is incorporated into the embryo as bowel.
- Midgut remains open.

Cephalocaudal and lateral folding occur simultaneously




- Meeting and fusion of cranial, lateral, and caudal edges of the embryo create the primordial foregut and hindgut
 - Slow fusion of midgut-due to presence of yolk sac. Midgut remains open until week 6-connects to yolk sac via *vitelline duct*.
 - Buccopharyngeal membrane opens at 4 and cloacal membrane at 7 weeks


Flexion delimits the bowel



- After the gut forms, it is attached to the body wall by dorsal and ventral mesenteries; ventral is lost except in region of liver. Vertebral duct remains in umbilical cord.



Anterior-posterior and lateral folding form the primitive gut



- Embryonic disc grows faster in length than the yolk sac causing the embryo to bend.
 - Dorsal surface grows more rapidly than the ventral
- Lateral folding
 - Fusion with apposing side except in the region of the yolk sac, and allantois
- Folding brings the heart and septum transversum caudal to bucco-pharyngeal membrane.




Figure 6–3. Formation of the dorsal mesentery. *A*, The primitive gut tube initially hangs from the posterior body wall by a broad bar of mesenchyme but, *B*, in regions inferior to the septum transversum this connection thins out to form a membranous dorsal mesentery composed of reflected peritoneum.

The dorsal mesentery thins to allow the gut to be flexibly suspended

The foregut has many derivatives


- Pharynx and its derivatives
- Lower Respiratory tract
- Esophagus
- Stomach
- Duodenum proximal to ampulla of Vater
- Liver
- Biliary Apparatus
- Pancreas

From stomach to biliary apparatus, all are supplied by the celiac artery, “the artery of the foregut.”


Esophagus elongates rapidly

- Appears to grow faster at its cranial than caudal end.
- Stomach does not descend but arises from a region just caudal to septum transversum that has been fated to be stomach.
- Epithelium obliterates lumen of esophagus and is recanalized by apoptosis (week 8).
 - Failure causes polyhydramnios
 - Esophageal atresia or tracheo-esophageal fistula.
- Stomach enlarges and rotates


Obliteration of the lumen and recanalization occurs



The stomach rotates 90° in a clockwise direction



- Dorsal surface grows faster than the ventral to create the greater and lesser curvature. Acquires a transverse position



Rotation of the stomach creates the lesser sac

- Dorsal mesogastrium moves to left.
- Ventral mesogastrium attaches to liver and body wall.
- Inferior recess form the greater omentum
 - Layers fuse to obliterate the lesser sac

**Rotation of the stomach
forms the omental bursa**



Movements of the mesentery and stomach are made possible by vacuolization due to selective apoptosis






Figure 9-4. The rotation of the stomach around its longitudinal axis commences with vacuolization of the right side of the thick mesenchymal bar that initially suspends the stomach from the posterior body wall.

Liver, biliary system and pancreas arise from the duodenum



Hepatic diverticulum grows from the duodenum into the ventral mesentery

- 
- Begins ~ week 4
 - Divides into cranial and caudal buds.
 - Cranial bud grows faster and becomes the hepatic parenchyma;
 - Hematopoietic colonists arrive ~ week 6
 - Caudal bud gives rise to the biliary system.

Ventral mesentery forms falciform ligament, hepatic peritoneum, and lesser omentum





Figure 9–8. Formation of the liver and associated membranes. As the liver bud grows into the ventral mesentery, its expanding crown makes direct contact with the developing diaphragm. The ventral mesentery that encloses the growing liver bud differentiates into the visceral peritoneum of the liver, which is reflected onto the diaphragm. This zone of reflection, which encircles the area where the liver directly contacts the diaphragm (the bare area), becomes the coronary ligament. The remnant of ventral mesentery connecting the liver with the anterior body wall becomes the falciform ligament, whereas the ventral mesentery between the liver and lesser curvature of the stomach forms the lesser omentum.

Ventral mesogastrium supports liver and stomach



Rotation of the stomach shapes the pancreas



- **Pancreas arises from dorsal and ventral buds.**
- **Rotation brings ventral to dorsal bud.**
- **Buds fuse.**
 - **Ventral duct becomes the main pancreatic duct but the dorsal bud forms most of the pancreas**
 - **Ventral bud forms only the uncinete process and inferior part of the head of the pancreas.**

Aberrant rotation causes an annular pancreas






Figure 9-7. The ventral pancreas may consist of two lobes. If the lobes migrate around the duodenum in opposite directions to fuse with the dorsal pancreatic bud, an annular pancreas is formed.




Derivatives of the midgut


- Small intestine (except for the proximal duodenum).
 - Cecum
 - Appendix
 - Ascending colon
 - Right 1/2 to 2/3 of the proximal transverse colon
- All are supplied by the superior mesenteric artery (“the artery of the midgut”)



**The midgut
grows
rapidly and
herniates
into the
umbilical
cord**

Week 6





**The midgut
rotates
around an
axis of the
superior
mesenteric
artery:**

1. 90°


2. 180°

Midgut hernia reduced at
week 10.

Rotation of the midgut

- 1. Cranial and caudal loop form.
- 2. Cranial growth >>> caudal growth.
- 3. Apex of loop is vitelline duct.
- 4. Cranial loop moves to right and caudal loop to left (90° counterclockwise).
- 4. Reduction of midgut hernia with rotation a further 180°.
 - Brings cecum to right
 - Moves down
 - Becomes secondarily retroperitoneal.

Loops of bowel fuse with the body wall and become secondarily retroperitoneal








Figure 9-10. Intraperitoneal, retroperitoneal, and secondarily retroperitoneal organs of the abdominal gastrointestinal tract.

**Volvulus is a serious
complication of excessive
flexibility**







A Persistent attachment to umbilicus Meckel's diverticulum



B Omphalomesenteric fistula



C Omphalomesenteric cyst




D Omphalomesenteric ligament (fibrous band)

Figure 9-21. Meckel's diverticulum. *A*, A typical Meckel's diverticulum is a finger-like projection of the ileum located about 100 cm proximal to the cecum. A Meckel's diverticulum may form (*B*) a patent fistula connecting the umbilicus with the ileum, (*C*) an isolated cyst suspended by ligaments, or (*D*) a fibrous band connecting the ileum and anterior body wall at the level of the umbilicus. (Drawings courtesy of Children's Hospital Medical Center, Cincinnati, OH.)


Derivatives of the hindgut


- Left 1/3 to 1/2 of the distal transverse colon
 - Descending colon
 - Sigmoid colon
 - Rectum
 - Superior part of anal canal
 - Epithelium of urinary bladder and most of the urethra
- **All are supplied by the inferior mesenteric artery, “the artery of the”. hindgut**

The hindgut is originally a cloaca-partioned to form rectum and urogenital sinus



Urorectal septum divides the cloaca





**Hindgut
forms
superior 2/3
of rectal
canal;
proctodeum
forms lower
1/3; divided
at pectinate
line**




**Never forget
the pectinate
line**


If anything can go wrong it will; anorectal malformations



The END

Have a nice day!





Timeline. Development of the gut tube and its derivatives.




Figure 9-22. Fate of the allantois. *A*, Normally, the allantois becomes occluded to form the urachus or median umbilical ligament of the adult. Very rarely, parts of the allantois may remain patent, producing (*B*) a urachal fistula, (*C*) a urachal sinus, (*D*) a urachal diverticulum, or (*E*) a urachal cyst.

Table 9-2

Derivatives of the Septum Transversum

| REGIONS OF SEPTUM TRANSVERSUM | DERIVATIVES |
|-----------------------------------|---|
| Cranial region | Central tendon of the diaphragm Myocytes of the pleuroperitoneal membranes |
| Central mesenchyme | Hematopoietic cells of liver |
| Caudal region (ventral mesentery) | Falciform ligament Visceral peritoneum of the liver, including the coronary ligament Visceral peritoneum of the gallbladder Lesser omentum, including the hepatoduodenal and hepatogastric ligaments |

Table 9-1

The Derivatives of the Primitive Gut Tube

| REGIONS OF THE DIFFERENTIATED GUT TUBE | ACCESSORY ORGANS DERIVED FROM THE GUT TUBE ENDODERM |
|---|--|
| <i>Foregut</i> | |
| Pharynx | Pharyngeal pouch derivatives (see Ch. 12) |
| Thoracic esophagus | Lungs |
| Abdominal esophagus | |
| Stomach | |
| Superior half of duodenum (superior to the ampulla of Vater) | Liver parenchyma and hepatic duct epithelium Gallbladder, cystic duct, and common bile duct Dorsal and ventral pancreatic buds (exocrine cells and pancreatic duct epithelium; probably also pancreatic endocrine cells) |
| <i>Midgut</i> | |
| Inferior half of duodenum | |
| Jejunum | |
| Ileum | |
| Cecum | |
| Appendix | |
| Ascending colon | |
| Right two thirds of transverse colon | |
| <i>Hindgut</i> | |
| Left one third of transverse colon | |
| Descending colon | |
| Sigmoid colon | |
| Rectum | Urogenital sinus and derivatives (see Ch. 10) |





Figure 9-1. A, The foregut, midgut, and hindgut of the primitive gut tube are formed by the combined action of differential growth and lateral and cephalocaudal folding. The foregut and hindgut are blind-ending tubes that terminate at the buccopharyngeal and cloacal membranes, respectively. The midgut is at first completely open to the cavity of the yolk sac B, C, As folding proceeds, however, this connection is constricted to form the narrow vitelline duct (D).