

## Facial and palatal development

L.Moss-Salentijn

---

---

---


---

---

---

---

## Timeline for development



---

---

---

---

---

---

---

## Contributions to the external face

- **Periprosencephalon:** ectoderm and mostly neural-derived mesenchyme surrounding the forebrain. Frontonasal process.
- **First pharyngeal (mandibular) arch.** Mandibular and maxillary processes.



---

---

---


---

---

---

---

## Contributions to external face




---

---

---


---

---

---

---

---




---

---

---

---


---

---

---

---

## Oropharyngeal membrane (buccopharyngeal, oral)



Membrane is composed of ectoderm and endoderm

---

---

---

---


---

---

---

---

## Disintegration of oropharyngeal membrane



Communication between foregut and amniotic cavity at approximately 4 weeks of development

---

---

---

---


---

---

---

---

## Stomodeum at 4 weeks



---

---

---

---

---

---

---

---

## Facial processes (prominences)



- Bilaterally:**  
Lateral nasal  
Medial nasal  
Maxillary  
Mandibular

---

---

---

---

---

---

---

---

**Development external face (4-5 wks)**



---

---

---

---


---

---

---

---

**Development external face (6-8 wks)**



---

---

---

---

---

---

---

---

**Face development animation 1**



---

---

---

---


---

---

---

---

**Dimensional changes (4-6 wks)**



---

---

---

---


---

---

---

---

**10-fold linear increase in size !**



---

---

---

---

---

---

---

---

**Face development – animation 2**



---

---

---

---

---


---

---

---

## Merging

**Differential mesenchymal proliferation. Elimination of groove.**



---

---

---

---


---

---

---


---

## **Merging with epithelial inclusion**



May result in facial cleft.

May be normal between LNP and maxillary process where enclosed epithelium gives rise to part of nasolacrimal duct epithelium.



---

---

---

---

---

---

---

---

## **Nasolacrimal duct between maxillary and lateral nasal processes**



---

---

---

---

---

---

---

---

### Sites of potential facial clefts



---

---

---

---

---

---

---

---

### Fusion

Contact and fusion of epithelium-covered surfaces. Removal of epithelium



---

---

---

---


---

---

---

---

### Fusion in primary and secondary palate development



---

---

---

---

---

---

---

---

## Fate of fused epithelium

- Non-proliferating epithelium in rapidly growing environment: passive stretch and incorporation in nearby surface epithelia
- Apoptosis and phagocytosis
- Epithelial-mesenchymal transformation

---

---

---

---


---

---

---

---

## Development of nose



---

---

---


---

---


---

---

---



Initial fusion of medial and lateral nasal processes, and subsequently between medial nasal and maxillary processes.



---

---

---


---

---

---

---

---




---



---



---




---



---



---




---



---



---




---



---



---




---



---



---



---



---




---



**Oronasal membrane**

Breaks down at about 6 wks of development.




---

---

---

---


---

---

---

---

**Primary (primitive) palate**



Primary palate composed of: intermaxillary segment of merged MNP's and the rostral tips of the maxillary processes. P: primary (primitive) choana permitting oro-nasal communication

---

---

---

---


---

---

---

---

**Development of primary and secondary palate**



Secondary palate development

---

---

---

---

---

---

---

---

**Intrinsic factors in the successful development of the secondary palate: increase in size of palatal processes**

- Mesenchymal cell proliferation – ceases hours before palatal processes become horizontal
- ECM production increasing volume of palatal processes
- Hydration of ECM – major increase in volume and turgor just prior to horizontalization

---

---

---

---


---

---

---

---

**Secondary palate development**



Palatal processes develop on the oral surfaces of the maxillary processes: initially vertically oriented, they assume horizontal orientation during eighth week of development.

---

---

---

---

---

---

---

---

**Horizontalization of palatal processes**



---

---

---

---

---

---

---

---

**Factors contributing to the horizontalization of the palatal processes**

- Turgor in the palatal processes
- Movements of the tongue – primitive swallowing- allowing tongue to move out of the way
- Downward and forward growth of lower jaw complex – providing space for the secondary palate
- Straightening of the cranial base – providing mechanical conditions for horizontalization

---

---

---


---

---

---

---

---



---

---

---

---

---

---

---

---



---

---

---

---

---

---

---

---

**Factors contributing to the successful fusion of the secondary palate: the medial edge epithelium (MEE)**

- Apoptosis of MEE surface cells immediately prior to fusion
- Development of temporary glycoprotein membrane coating, enabling adhesion between MEE cells of opposing palatal processes
- Successful removal of MEE from fusion line

---

---

---


---

---

---

---

---



Fate of MEE cells: apoptosis (TUNEL reaction above) and phagocytosis

---

---

---


---

---

---

---

---



Non-proliferating epithelium in rapidly growing environment: passive stretch and incorporation in nearby surface epithelia

---

---

---

---


---

---

---

---

### Completion of palate formation



---

---

---

---


---

---

---

---

### Sites of potential palatal clefts



---

---

---

---

---

---

---

---

### PLACODES

Localized thickened areas of specialized ectoderm, lateral to the neural crest, at the border between neural plate and the future epidermis.

---

---

---


---

---

---

---

---




---

---

---

---

---


---

---

---

### Location of placodes

- *Near forebrain :*
  - ◆ Olfactory placode
  - ◆ Lens placode




---

---

---

---

---


---

---

---

### Location of placodes

- *Dorsolateral :*
  - ◆ Otic placode: related to (= evolved from or having common origin with) lateral line system




---

---

---

---

---

---

---

---

## Development of organs of special sense

Surface ectoderm	Nervous System	ORGAN OF SENSORY ELEMENT	ORGAN OF NERVOUS CONDUCTION
Gen. epithelial cell	INTEL. CNS	FOVEAE	FOVEAE
	VISION	NEURAL TUBE	NEURAL TUBE
	AUDITION & BALANCE	FOVEAE	FOVEAE
	TASTE	Sensory differentiation of surface cells of surface ectodermal covering of tongue	Neural crest Nerve ganglion
L. Epithelial cells M. Mesoderm	TRIGEM. SENSORY	Free nerve endings to nasal and mechanical sense, etc.	Neural crest Nerve ganglion

---

---

---

---

---


---

---

---

---

---




---

---

---

---

---


---

---

---

---

---




---

---

---

---

---


---

---

---

---

---




---



---



---



---




---



---

## Branchiomic nerves: origins and axon projection patterns




---



---



---



---



---



---