


# Entodermal derivatives: formation of the gut, liver, and pancreas


Mike Gershon



## Folding forms the gut


- Primitive gut extends from buccopharyngeal to cloacal membrane.
  - Move toward each other
- Cardiogenic mesenchyme is originally rostral, but folding brings it caudal to buccal membrane.
- Foregut and hindgut become recognizable
- Portion of yolk sac is incorporated into the embryo as bowel.
- Midgut remains open.

## Cephalocaudal and lateral folding occur simultaneously




- Meeting and fusion of cranial, lateral, and caudal edges of the embryo create the primordial foregut and hindgut
  - Slow fusion of midgut-due to presence of yolk sac. Midgut remains open until week 6-connects to yolk sac via vitelline duct.
  - Buccopharyngeal membrane opens at 4 and cloacal membrane at 7 weeks

## Flexion delimits the bowel




- After the gut folds, the dorsal and ventral walls meet except in the region of the liver and stomach.

## Anterior-posterior and lateral folding form the primitive gut



- Embryonic disc grows faster in length than the yolk sac causing the embryo to bend.
  - Dorsal surface grows more rapidly than the ventral
- Lateral folding
  - Fusion with apposing side except in the region of the yolk sac, and allantois
- Folding brings the heart and septum transversum caudal to buccopharyngeal membrane.



## The dorsal mesentery thins to allow the gut to be flexibly suspended

Figure 6-3. Formation of the dorsal mesentery. A, The primitive gut tube initially hangs from the posterior body wall by a broad bar of mesenchyme but, B, in regions inferior to the septum transversum this connection thins out to form a membranous dorsal mesentery composed of reflected peritoneum.


## The foregut has many derivatives

- Pharynx and its derivatives
- Lower Respiratory tract
- Esophagus
- Stomach
- Duodenum proximal to ampulla of Vater
- Liver
- Biliary Apparatus
- Pancreas

From stomach to biliary apparatus, all are supplied by the celiac artery, "the artery of the foregut."

## Esophagus elongates rapidly

- Appears to grow faster at its cranial than caudal end.
- Stomach does not descend but arises from a region just caudal to septum transversum that has been fated to be stomach.
- Epithelium obliterates lumen of esophagus and is recanalized by apoptosis (week 8).
  - Failure causes polyhydramnios
  - Esophageal atresia or tracheo-esophageal fistula.
- Stomach enlarges and rotates



## Obliteration of the lumen and recanalization occurs




Figure 9-13. Formation of the definitive gut lumen. Proliferation of the endodermal lining completely occludes the gut tube during the sixth week. Recanalization is completed by week 9. Incomplete or abnormal recanalization may result in duplication of the lumen or stenosis of the gut tube.

## The stomach rotates 90° in a clockwise direction





Figure 9-3. Rotations of the stomach. A-C, Oblique frontal views; D, direct frontal view. The posterior wall of the stomach expands during the fourth and fifth weeks to form the greater curvature. During the seventh week, the stomach rotates clockwise on its longitudinal axis (when viewed from above).


- Dorsal surface grows faster than the ventral to create the greater and lesser curvature. Acquires a transverse position

## Rotation of the stomach creates the lesser sac


- Dorsal mesogastrum moves to left.
- Ventral mesogastrum attaches to liver and body wall.
- Inferior recess forms the greater omentum
  - Layers fuse to obliterate the lesser sac




## Rotation of the stomach forms the omental bursa




## Movements of the mesentery and stomach are made possible by vacuolization due to selective apoptosis




## Liver, biliary system and pancreas arise from the duodenum




## Hepatic diverticulum grows from the duodenum into the ventral mesentery




## Ventral mesentery forms falciform ligament, hepatic peritoneum, and lesser omentum



## Ventral mesogastrium supports liver and stomach



## Rotation of the stomach shapes the pancreas



## Aberrant rotation causes an annular pancreas




Figure 9-7. The ventral pancreas may consist of two lobes. If the lobes migrate around the duodenum in opposite directions to fuse with the dorsal pancreatic bud, an annular pancreas is formed.




Figure 9-2. Structure of the gut tube. The foregut consists of the pharynx, located superior to the respiratory diverticulum, the thoracic esophagus, and the abdominal foregut. The abdominal foregut forms the abdominal esophagus, stomach, and about half of the duodenum and gives rise to the liver, the gallbladder, the pancreas, and their associated ducts. The midgut forms half of the duodenum, the jejunum and ileum, the ascending colon, and about two thirds of the transverse colon. The hindgut forms one third of the transverse colon, the descending colon and sigmoid colon, and the upper two thirds of the anorectal canal. The abdominal esophagus, stomach, and superior part of the duodenum are suspended by dorsal and ventral mesenteries; the abdominal gut tube excluding the rectum is suspended in the abdominal cavity by a dorsal mesentery only.

## Derivatives of the midgut

- Small intestine (except for the proximal duodenum).
- Cecum
- Appendix
- Ascending colon
- Right 1/2 to 2/3 of the proximal transverse colon
- All are supplied by the superior mesenteric artery ("the artery of the midgut")




Figure 9-3. Development of the greater omentum and lesser sac. A, B. The rotation of the stomach and stomach of the dorsal mesogastrium results in a sac (the greater omentum) that hangs from the greater curvature of the stomach. The lesser mesogastrium migrates to the right, becomes posteriorly fixed to the body wall, enclosing the space posterior to the stomach and within the expanding cavity of the greater omentum. This space is the lesser sac of the peritoneal cavity. The remainder of the peritoneal cavity is now called the greater sac. The principal passageway between the greater and lesser sacs is the epiploic foramen of Winslow.

The midgut grows rapidly and herniates into the umbilical cord

Week 6





Figure 9-6. Migration and rotation of the intestines. A, B. At the end of the sixth week, the primary foregut has herniated into the umbilical cord through the epigastric foramen of the body wall. C. The small intestine elongates to form four-and-a-half loops, the cecum and appendix grow, and at the end of the 10th week, the primary foregut body retracts into the abdominal cavity, creating an additional half loop (the lesser sac) posterior to the stomach. The stomach is now displaced inferiorly, giving greater the potential mobility for such an expanding sac. The ascending colon is retroperitoneal from the side of the posterior abdominal wall. The pylorus, duodenum, and transverse colon and sigmoid colon remain suspended by mesentery.

The midgut rotates around an axis of the superior mesenteric artery:

1. 90°
2. 180°

Midgut hernia reduced at week 10.

## Rotation of the midgut

- 1. Cranial and caudal loop form.
- 2. Cranial growth >>> caudal growth.
- 3. Apex of loop is vitelline duct.
- 4. Cranial loop moves to right and caudal loop to left (90° counterclockwise).
- 4. Reduction of midgut hernia with rotation a further 180°.
  - Brings cecum to right
  - Moves down
  - Becomes secondarily retroperitoneal.

## Loops of bowel fuse with the body wall and become secondarily retroperitoneal




Figure 6-4. The distinction between intraperitoneal, retroperitoneal and secondarily retroperitoneal positions of the viscera. A Viscera suspended within the peritoneal cavity by a mesentery are called intraperitoneal, whereas organs embedded in the body wall and covered by peritoneum are called retroperitoneal. B The mesentery suspending some intraperitoneal organs disappears as both mesentery and organ fuse with the body wall. These organs are then called secondarily retroperitoneal.




Figure 9-10. Intraperitoneal, retroperitoneal, and secondarily retroperitoneal organs of the abdominal gastrointestinal tract.

## Volvulus is a serious complication of excessive flexibility




Figure 9-20. Volvulus: Volvulus may occur as suspended regions of the gut twist around themselves, constricting the intestine and/or compromising its blood supply.





Figure 9-21. Meckel's diverticulum. A, A typical Meckel's diverticulum is a finger-like projection of the ileum located about 100 cm proximal to the cecum. A Meckel's diverticulum may form (B) a patent fistula connecting the ileum with the umbilicus, (C) an isolated cyst suspended by ligaments, or (D) a fibrous band connecting the ileum and anterior body wall at the level of the umbilicus. (Drawings courtesy of Children's Hospital Medical Center, Cincinnati, OH.)

## Derivatives of the hindgut

- Left 1/3 to 1/2 of the distal transverse colon
- Descending colon
- Sigmoid colon
- Rectum
- Superior part of anal canal
- Epithelium of urinary bladder and most of the urethra
- **All are supplied by the inferior mesenteric artery, "the artery of the" hindgut**

## The hindgut is originally a cloaca-partioned to form rectum and urogenital sinus



## Urorectal septum divides the cloaca




Figure 9-11. Subdivision of the cloaca into an anterior primitive urogenital sinus and a posterior rectum between 4 and 6 weeks. The urorectal septum that divides the cloaca is composed of three distinct septa. Initially, a superior Tourneux fold grows inferiorly to the level of the future pelvic urethra. Separation is then completed by left and right Rathke folds that grow in a coronal plane.







Figure 9-12. The lower third of the ano-rectal canal is formed by an ectodermal invagination called the anal pit. The border between the superior end of the anal pit and the inferior end of the rectum is demarcated by mucosal folds called the pectinate line in the adult.


Hindgut forms superior 2/3 of rectal canal; proctodeum forms lower 1/3; divided at pectinate line



Never forget the pectinate line

## If anything can go wrong it will; anorectal malformations






**Table 9-2**  
**Derivatives of the Septum Transversum**

REGIONS OF SEPTUM TRANSVERSUM	DERIVATIVES
Cranial region	Central tendon of the diaphragm Myocytes of the pleuroperitoneal membranes
Central mesenchyme	Hematopoietic cells of liver
Caudal region (ventral mesentery)	Falciform ligament Visceral peritoneum of the liver, including the coronary ligament Visceral peritoneum of the gallbladder Lesser omentum, including the hepatoduodenal and hepatogastric ligaments

**Table 9-1**  
**The Derivatives of the Primitive Gut Tube**

REGIONS OF THE DIFFERENTIATED GUT TUBE	ACCESSORY ORGANS DERIVED FROM THE GUT TUBE ENDODERM
<b>Foregut</b> Pharynx Thoracic esophagus Abdominal esophagus Stomach Superior half of duodenum (superior to the ampulla of Vater)	Pharyngeal pouch derivatives (see Ch. 12) Lungs
<b>Midgut</b> Inferior half of duodenum Jejunum Ileum Cecum Appendix Ascending colon Right two thirds of transverse colon	Liver parenchyma and hepatic duct epithelium Gallbladder, cystic duct, and common bile duct Dorsal and ventral pancreatic buds (exocrine cells and pancreatic duct epithelium; probably also pancreatic endocrine cells)
<b>Hindgut</b> Left one third of transverse colon Descending colon Sigmoid colon Rectum	Urogenital sinus and derivatives (see Ch. 10)



**The END**

Have a nice day!