


HD9 - Development of the Kidney and Ureter




nephrons in the kidney generate urine that is propelled to the ureters and then to the bladder for storage and excretion


The Urinary outflow tract:

- ◆ monitors and regulates extra-cellular fluids
- ◆ excretes harmful substances in urine, including nitrogenous wastes (urea)
- ◆ returns useful substances to bloodstream
- ◆ maintain balance of water, electrolytes (salts), acids, and pH in the body fluids


The urogenital system derives predominantly from intermediate mesoderm




During development, 3 successive kidneys form:



- Pronephros (head kidney)
- Mesonephros (middle kidney)
- Metanephros (definitive kidney)



pronephros in an early embryo



Mesonephros in intermediate embryo

HD9 - Development of the Kidney and Ureter

C. Metanephros in late embryo and adult

A **metanephros** is always drained exclusively by one duct, the ureter.
 In birds in reptiles the ureter separates from the **nephric duct (Wolffian duct)** and enters the **cloaca**. In mammals, the ureter separates from the nephric duct and enters the bladder

A. Early
 B. Late

renal development begins when the **ureteric bud** invades kidney mesenchyme (**the metanephric blastema**)

As the embryo grows, the **ureters lengthen**, and the **kidneys rotate** and **ascend** along the dorsal body wall

The kidney is radially patterned

- branching morphogenesis and nephron formation last until just after birth
- occur exclusively in the peripheral domain beneath the renal capsule
- new generations of nephrons and ureter branches displace older generations inward
- further differentiation occurs in inner domains at a distance from the renal capsule

RECIPROCAL SIGNALING BETWEEN STROMA, NEPHRON PROGENITORS AND URETERIC BUD TIPS GIVES RISE TO CELL TYPES IN THE MATURE KIDNEY


nephron progenitors	NEPHRONS
ureteric bud tips	COLLECTING DUCT SYSTEM
stroma	CAPSULE/INTERSTITIUM

Induction → MET → Branching

Labels: Condensed mesenchyme, Ureteric bud tips, Renal vesicle, Stroma, S-shaped body, Nephron, Distal tubule, Proximal tubule, Glomerulus, Endothelial cells, Collecting duct, Efferent arteriole, Afferent arteriole, Podocytes, Peritubular capillary endothelium, Thick ascending limb, Henle's loop, Proximal tubule, Distal tubule, Collecting duct.

HD9 - Development of the Kidney and Ureter


the collecting duct system and ureter are derived from the ureteric bud



branching ureteric bud tips
collecting ducts
ureter

The distinct cellularity of the collecting duct system and ureter depends on developmental signals from surrounding mesenchyme

shape changes and local proliferation at ureteric bud tips forms an **ampulla**





Branching morphogenesis:

- ampullae form at ureteric bud tips
- a cleft forms and the tips begin to bifurcate
- the tips elongate
- new ampullae form


day 1
Wolffian duct
ub
2ndG
3rdG
4thG

The collecting duct system grows from the periphery by **dichotomous branching**

at birth:


Branching morphogenesis in Real time



00:00:00


Costantini Lab
Columbia University, Dept. of Genetics & Development

NEPHRONS FORM EXCLUSIVELY AT **URETERIC BUD TIPS** IN RESPONSE TO LOCAL SIGNALS FROM URETERIC BUD CELLS



nephron progenitors
ureteric bud tip

Nephron progenitors condense at ub tips, **aggregate**



and **trans-differentiate** into epithelial cells that make up the renal vesicle, **Comma** and **S-shaped bodies**

from: The kidney, Eds. Vize et al., 2003

TRENDS in Cell Biology

HD9 - Development of the Kidney and Ureter

RET-Gdnf signaling is crucial for branching morphogenesis

GDNF signals through GFRa-1 and RET

Ret mutations in humans cause renal abnormalities, Hirschsprung's disease and cancer

GDNF signals through GFRa-1 and RET

RET TYROSINE KINASE

URETERIC BUD FORMATION

BRANCHING MORPHOGENESIS

RENAL AGENESIS

URETER AGENESIS

RENAL HYPODISPLASIA

Mutations in Ret, Gdnf or Gfra1 result in renal agenesis or hypoplasia

STROMAL CELL SECRETE RETINOIC ACID, THE ACTIVE FORM OF VITAMIN A
VITAMIN A INDUCES RET EXPRESSION IN NEARBY URETERIC BUD CELLS

STROMA

URETERIC BUD

RA

NEPHRON PROGENITORS

vitamin A

Ret/Gfra1

Gdnf

Gdnf secreted by nephron progenitors binds to Ret via the Ret co-receptor (Gfra1) inducing branching morphogenesis

The Ret receptor is expressed in ureteric bud tips and controls branching morphogenesis

ureteric bud (RET)

stroma (vitamin A)

ub

Vitamin A from Stromal cells controls Ret expression in ureteric bud cells

Vitamin A deficiency generates renal malformations similar to those induced by Ret mutations

ureteric bud cells must express Ret to contribute to a tip

a

Hoxb7/GFP, Ret^{-/-} ES cells

wild type blastocyst

chimeric embryo

b

UB

WD

c

d





e

control


What cellular processes are stimulated by GDNF to promote branching of the ureteric epithelium? Some candidates are:

- localized cell proliferation
- changes in cell shape
- migration
- differential cell-cell adhesion
- localized remodeling of the extracellular matrix


HD9 - Development of the Kidney and Ureter



	<p><u>abnormal position of the ureter orifice</u> vitamin A deficiency, Ret sprouty, slit-2, retinoid excess</p>
<p>Physical vs Functional obstruction</p>	
	<p><u>abnormal peristalsis</u> sonic hedgehog (muscle) Calcineurin B (peristalsis) uroplakin (epithelium) Tbx18?</p>




HD9 - Development of the Kidney and Ureter



- The **urogenital sinus** forms the **bladder, urethra** (including the **prostate and penis**)
- The **mesonephric duct** (aka **Wolffian duct**) forms the **vas (ductus) deferens, seminal vesicle and epididymis** in males
- Mullerian ducts (paramesonephric ducts)** degenerate in males


from: The kidney: Eds, Vize et al., 2003)



- in females the **urogenital sinus** forms the **bladder, urethra and vagina**
- Mullerian (paramesonephric ducts)** differentiate into the **uterus and upper vagina**
- Wolffian (mesonephric ducts)** regress


from: The kidney: Eds, Vize et al., 2003)

The trigone is the center of the anti-reflux mechanism




The trigone is bounded by the ureter orifices at its lateral edges and the sex ducts, at the apex

Abnormal connections between the ureter orifice and trigone are associated with vesicoureteral reflux and obstruction





Mackie-Stephens hypothesis: the final position of the ureter with respect to the Trigone depends on the site of its formation on the Wolffian duct

The trigone is a region where the detrusor and urethral muscle join the ureteral fibers









proper configuration of muscle groups that form the trigone is likely to be important for urinary tract function

the **ureteral valve** is part of the trigone and is an **anti-reflux** mechanism that prevents urine back flow (reflux)




Ureteral valve function depends on insertion of the ureter orifice at the proper position in the bladder neck (trigone)

HD9 - Development of the Kidney and Ureter




HD9 - Development of the Kidney and Ureter

How do ureters move from the Wolffian duct to the bladder?



According to the accepted model, trigone formation is considered to be crucial for repositioning the ureter orifice



during ureter transposition, the **cnd** is incorporated into the bladder where it expands to form the **trigone** effectively separating the ureter orifice from the Wolffian duct


Accepted model of ureter transposition



formation of the **trigone** from the **common nephric duct** repositions the ureters in the bladder


Larsen's Embryology

using mouse models to re-assess the mechanism of ureter transposition:





expression of Jelly Fish green fluorescent protein in the mouse common nephric duct of this transgenic mouse enables us to follow its fate during ureter insertion

what happens to the common nephric duct during ureter transposition?



The common nephric duct appears to regress rather than expand

Ureter transposition depends on apoptosis of the **common nephric duct**



HD9 - Development of the Kidney and Ureter

