


Entodermal derivatives:
formation of the gut, liver,
and pancreas



Folding forms the gut


- Primitive gut extends from buccopharyngeal to cloacal membrane.
 - Move toward each other
- Cardiogenic mesenchyme is originally rostral, but folding brings it caudal to buccal membrane.
- Foregut and hindgut become recognizable
- Portion of yolk sac is incorporated into the embryo as bowel.
- Midgut remains open.

Cephalocaudal and lateral folding occur simultaneously

- Meeting and fusion of cranial, lateral, and caudal edges of the embryo create the primordial foregut and hindgut
 - Slow fusion of midgut-due to presence of yolk sac. Midgut remains open until week 6-connects to yolk sac via *vitelline duct*.
 - Buccopharyngeal membrane opens at 4 and cloacal membrane at 7 weeks

Flexion delimits the bowel


- After the gut forms, it is attached to the body wall by dorsal and ventral mesenteries; ventral is lost except in region of liver. Vitelline duct remains in umbilical cord.



Anterior-posterior and lateral folding form the primitive gut


- Embryonic disc grows faster in length than the yolk sac causing the embryo to bend.
 - Dorsal surface grows more rapidly than the ventral
- Lateral folding
 - Fusion with adjoining side except in the region of the yolk sac, and allantois
- Folding brings the heart and septum transversum caudal to buccopharyngeal membrane.

The septum transversum partially separates the thoracic and abdominal cavities




- The superior portion is the primitive pericardial cavity.
- The inferior portion is the future peritoneal cavity.
- The pericardial and peritoneal cavities communicate through the pericardioperitoneal canals

The pericardioperitoneal canals are closed by the formation of the pleuroperitoneal membranes




- The pleuroperitoneal membranes contribute muscle to the definitive diaphragm

The definitive diaphragm is a composite structure



1. Septum transversum
2. Pleuroperitoneal membranes
3. Paraxial mesoderm
4. Esophageal mesenchyme




The dorsal mesentery thins to allow the gut to be flexibly suspended

Figure 6-3. Formation of the dorsal mesentery. A, The primitive gut tube initially hangs from the posterior body wall by a broad bar of mesenchyme but, B, in regions inferior to the septum transversum this connection thins out to form a membranous dorsal mesentery composed of reflected peritoneum.

The gut is regionally specified early in development

- Endoderm is specified before gut tube is complete.
- Specification is manifest as a series of regionally specific transcription factors.
- The boundaries between regions, however, are plastic and depend on interactions with mesoderm

Signals and transcription factors specify regionalization of the gut



Schneuwelt et al. Larval Human Embryology, 4th Edition. Copyright © 2005 by Churchill Livingstone, an imprint of Elsevier, Inc. All rights reserved.

The boundaries of GI regions are set by endodermal mesodermal cross-talk

(B)

Endoderm	Mesoderm
(Localized transcription factors in endoderm) → Sonic hedgehog in endoderm	Reception of Sonic hedgehog protein in mesoderm
↓	↓
Reception of paracrine signals by endoderm	Hox gene expression in mesoderm
↓	↓
Specification and differentiation of endoderm	Specification of mesoderm
↓	↓
Paracrine factors	BMPs, FGFs made by mesoderm
↓	↓
Differentiation of mesoderm	↓

- The language is paracrine secretion.
 - Secreted growth factors
 - Detected by appropriate receptors.
 - Coupled to transduction channels that affect transcription.
- Begins with Shh expression in posterior endoderm- spreads to whole gut
 - Induces series of Hox genes in mesoderm
 - Mesoderm then influences epithelial differentiation

Mesenchyme critically affects epithelial differentiation

(C)

- Wnt signaling specifies intestinal epithelium.
 - Wnt = intestine
 - No Wnt = stomach
- Mesenchyme of stomach expresses Barx1 and secretes Wnt inhibitors (SFRP1, 2).
- Mesenchyme of intestine secretes BMP4
 - Induces mesenchyme anterior to it to express Sox9 + Nkx-2
 - Become pyloric sphincter

The foregut has many derivatives

- Pharynx and its derivatives
- Lower Respiratory tract
- Esophagus
- Stomach
- Duodenum proximal to ampulla of Vater
- Liver
- Biliary Apparatus
- Pancreas

From stomach to biliary apparatus, all are supplied by the celiac artery, "the artery of the foregut."

Esophagus elongates rapidly

- Appears to grow faster at its cranial than caudal end.
- Stomach does not descend but arises from a region just caudal to septum transversum that has been fated to be stomach.
- Epithelium obliterates lumen of esophagus and is recanalized by apoptosis (week 8).
 - Failure causes polyhydramnios
 - Esophageal atresia or tracheo-esophageal fistula.
- Stomach enlarges and rotates

Obliteration of the lumen and recanalization occurs

Figure 9-13. Formation of the definitive gut lumen. Proliferation of the endodermal lining completely occludes the gut tube during the sixth week. Recanalization is completed by week 9. Incomplete or abnormal recanalization may result in duplication of the lumen or stenosis of the gut tube.

The stomach rotates 90° in a clockwise direction

Figure 9-3. Rotations of the stomach. A-C, Oblique frontal views; D, direct frontal view. The posterior wall of the stomach expands during the fourth and fifth weeks to form the greater curvature. During the seventh week, the stomach rotates clockwise on its longitudinal axis (when viewed from above).

- Dorsal surface grows faster than the ventral to create the greater and lesser curvature. Acquires a transverse position

Rotation of the stomach creates the lesser sac

- Dorsal mesogastrum moves to left.
- Ventral mesogastrum attaches to liver and body wall.
- Inferior recess forms the greater omentum
 - Layers fuse to obliterate the lesser sac

Rotation of the stomach forms the omental bursa

Movements of the mesentery and stomach are made possible by vacuolization due to selective apoptosis

Figure 9-4. The rotation of the stomach around its longitudinal axis commences with vacuolization of the right side of the thick mesenchymal bar that initially suspends the stomach from the posterior body wall.

Liver, biliary system and pancreas arise from the duodenum



Figure 9-6. Development of the liver, gallbladder, pancreas, and their duct systems from endodermal diverticula of the duodenum. The liver bud sprouts during the fourth week and expands in the ventral mesentery. The cystic diverticulum and ventral pancreatic bud also grow into the ventral mesentery, whereas the dorsal pancreatic bud grows into the dorsal mesentery. During the fifth week, the ventral pancreatic bud migrates around the posterior side (latter right side) of the duodenum to fuse with the dorsal pancreatic bud. The main duct of the ventral bud ultimately becomes the major pancreatic duct, which drains the entire pancreas.

Hepatic diverticulum grows from the duodenum into the ventral mesentery

- Begins ~ week 4
- Divides into cranial and caudal buds.
- Cranial bud grows faster and becomes the hepatic parenchyma;
 - Hematopoietic colonists arrive ~ week 6
- Caudal bud gives rise to the biliary system.




Ventral mesentery forms falciform ligament, hepatic peritoneum, and lesser omentum

Figure 9-8. Formation of the liver and associated membranes. As the liver bud grows into the ventral mesentery, its expanding crown makes direct contact with the developing diaphragm. The ventral mesentery that encloses the growing liver bud differentiates into the visceral peritoneum of the liver, which is reflected onto the diaphragm. This zone of reflection, which encircles the area where the liver directly contacts the diaphragm (the bare area), becomes the coronary ligament. The remnant of ventral mesentery connecting the liver with the anterior body wall becomes the falciform ligament, whereas the ventral mesentery between the liver and lesser curvature of the stomach forms the lesser omentum.




Pancreas arises from dorsal and ventral buds.

- Rotation brings ventral to dorsal bud.
- Buds fuse. Ventral duct becomes the main pancreatic duct but the dorsal bud forms most of the pancreas
 - Ventral bud forms only the uncinate process and inferior part of the head of the pancreas




Derivatives of the midgut

- Small intestine (except for the proximal duodenum).
- Cecum
- Appendix
- Ascending colon
- Right 1/2 to 2/3 of the proximal transverse colon
- All are supplied by the superior mesenteric artery ("the artery of the midgut")



The midgut grows rapidly and herniates into the umbilical cord


Week 6



The midgut rotates around an axis of the superior mesenteric artery

Rotation of the midgut

1. Cranial and caudal loop form.
2. Cranial growth >>> caudal growth.
3. Apex of loop is vitelline duct.
4. Cranial loop moves to right and caudal loop to left (90° counterclockwise).
4. Reduction of midgut hernia with rotation a further 180°.
 - Brings cecum to right
 - Moves down
 - Becomes secondarily retroperitoneal.



Loops of bowel fuse with the body wall and become secondarily retroperitoneal







Figure 9-10. Intraperitoneal, retroperitoneal, and secondarily retroperitoneal organs of the abdominal gastrointestinal tract.



- ### Derivatives of the hindgut
- Left 1/3 to 1/2 of the distal transverse colon
 - Descending colon
 - Sigmoid colon
 - Rectum
 - Superior part of anal canal
 - Epithelium of urinary bladder and most of urethra
 - **All are supplied by the inferior mesenteric artery, "the artery of the". hindgut**

